

Resonant Tunneling Between Quantum Hall Edge States

K. Moon[1], H. Yi[2], C.L. Kane[2], S.M. Girvin[1], Matthew P.A. Fisher[3]

[1] *Department of Physics, Indiana University, Bloomington, Indiana 47405*

[2] *Department of Physics, University of Pennsylvania, Philadelphia, PA 19104*

[3] *IBM Research, T.J. Watson Research Cent., PO Box 218, Yorktown Heights, NY 10598*

(February 1, 2008)

Abstract

Resonant tunneling between fractional quantum Hall edge states is studied in the Luttinger liquid picture. For the Laughlin parent states, the resonance line shape is a universal function whose width scales to zero at zero temperature. Extensive quantum Monte Carlo simulations are presented for $\nu = 1/3$ which confirm this picture and provide a parameter-free prediction for the line shape.

PACS numbers: 72.10.-d 73.20.Dx

Many body correlations often play an important role in tunneling and resonant tunneling in mesoscopic structures, such as quantum dots [1,2]. In addition to the Coulomb correlations in the vicinity of a tunneling structure, it has recently been emphasized that the electron interactions in the “leads” feeding the structure can be crucial. Specifically, when the leads are one-dimensional, interactions are believed to de-stabilize the Fermi liquid, forming a Luttinger liquid. In this case, the tunneling conductance through a point contact is predicted to vanish as a power law of temperature, due to the absence of long-lived single-particle excitations in the incident electron gas [3]. Unfortunately, in real one-dimensional wires extraneous impurities away from the point contact will complicate matters, tending to backscatter and localize the electrons.

In this article, we consider the situation where the “leads” feeding a point contact are 2d fractional quantum Hall fluids. We argue that the correlations built into Laughlin states play a crucial role in determining the nature of resonant tunneling at low temperatures. In particular, we show that as the temperature is lowered, resonances become sharper and have a universal shape. We present the results of extensive quantum Monte Carlo simulations which give the universal line shape expected for the $\nu = 1/3$ state.

Provided the temperature is below the gap in the fractional quantum Hall fluid, the incoming current will be carried by edge states. In a beautiful series of articles, Wen [4] has demonstrated that the gapless edge excitations of a quantum Hall system are “chiral” Luttinger liquids. Since the electrons in a chiral Luttinger liquid move only in one direction, backscattering is only possible when opposite edges of the sample are close together, namely at the point contact. Thus localization in the edge state “leads” is completely unimportant.

Recently, two of us have studied the problem of resonant tunneling in a Luttinger liquid [7]. A renormalization group analysis reveals that the resonance lineshape at low temperatures is universal, depending only on the two terminal conductance, $G = ge^2/h$ of the Luttinger liquid. The parameter g depends in a complicated way on the strength of the interactions between left- and right-moving electrons. In a quantum Hall edge state the electrons move only in one direction. Most strikingly, in this case g is a topological invari-

ant controlled by the quantum Hall state in the bulk [4]. To see this, consider raising the chemical potential of the right-movers relative to that of the left by an amount $\delta\mu$. This corresponds to applying a Hall voltage $V_H = \delta\mu/e$ and the resulting current is given by the quantized Hall coefficient as $I = \nu(e^2/h)V_H$. This immediately establishes the universal result within a given Hall plateau: $g = \nu$. *This remarkable fact makes the resonance line shape completely universal, model-independent and fully determined (up to an overall temperature scale).* The fractional quantum Hall regime ($g < 1$) is thus a far more promising place to observe pure Luttinger liquid behavior than in a single-channel quantum wire, where it is difficult to eliminate disorder and where the value of g is unknown. Chamon and Wen [5] have recently considered a theory of resonant tunneling between quantum Hall edge states which is valid in the limit in which the peak conductance of the resonances are much less than e^2/h , or in the tails of a stronger resonance. A similar theory for resonant tunneling in a Luttinger liquid has been developed by Furusaki and Nagaosa [6]. In contrast, the scaling theory presented below is valid over the entire width of the resonance for resonances whose peak conductance approaches the ‘perfect’ value $\nu e^2/h$.

The analog of a weak impurity that causes back-scattering is a narrow constriction which brings the left- and right-movers close enough together to communicate via tunneling of Laughlin quasiparticles through a ‘weak link’ as illustrated in Fig.(1). This can be achieved by means of a gate which electrostatically defines a narrow region in the Hall bar. The analog of the two-impurity resonance geometry considered by Kane and Fisher [7] would be two nearby, symmetric tunneling paths for quasiparticles [8]. For some value of magnetic field or gate voltage one will (randomly or intentionally) achieve the condition of destructive interference [9] which shuts off the inter-edge quasiparticle tunneling. This is the resonance (no backscattering) condition which will be manifested experimentally by the appearance of a two-terminal source-drain conductance [10] which peaks at a value which at low temperatures approaches the quantized value, $G = \nu e^2/h$. Away from resonance the quasi-particle tunneling causes current to leak from one edge to the other, thereby reducing the source to drain conductance (see Fig.(1). In fact, in the fractional quantum Hall effect the quasipar-

ticle tunneling is expected to increase upon cooling, driving the conductance all the way to zero in the zero temperature limit.

Following the analysis of ref. [7], the weak quasiparticle tunneling regime can be analyzed perturbatively. As the temperature is lowered and the effective tunneling amplitude grows, perturbative analysis fails. However at low enough temperatures the conductance from source to drain becomes tiny and a perturbative analysis in terms of electron transmission becomes possible. The peak and tail regimes of the resonance are tied together by a universal scaling function. Below we establish a model and calculate the scaling function for $g = 1/3$ using quantum Monte Carlo.

We begin our analysis by briefly reviewing the logic behind Wen's edge state theory. For simplicity we focus here on the primary Hall states with inverse filling factor ν^{-1} equal to an odd integer. In this case the edge state has only one branch.

Conservation of electron three-current j_μ permits us to introduce a fictitious gauge field a_μ via

$$j_\mu = \frac{1}{2\pi} \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda. \quad (1)$$

The bulk 2D electron gas is in an incompressible quantum Hall state with an excitation gap, which means that the low-energy, long-length-scale physics must be described by a massive theory. In 2+1-D the only massive gauge theory is the Chern-Simons theory which has (Euclidean) action [11] (ignoring irrelevant terms):

$$S_{bulk} = \frac{i}{4\pi\nu} \int a_\mu \partial_\nu a_\lambda \epsilon_{\mu\nu\lambda} d^2x d\tau, \quad (2)$$

The coefficient ν^{-1} is uniquely fixed by the quantized Hall conductivity and specifies the number of zeros bound to the electrons in the Laughlin wave function [12–14]

Wen has shown that in the presence of a boundary, say at $y=0$, an effective action for the edge state can be obtained as follows [4]: First integrate out a_τ in the bulk, which gives an incompressibility constraint on the electron density, $\epsilon_{ij} \partial_i a_j = 0$. Then solve the constraint in terms of a scalar field, $a_j = \partial_j \phi$. After an integration by parts the final Euclidean action for the edge state takes the form,

$$S_{\text{edge}} = \frac{1}{4\pi g} \int dx d\tau (\partial_x \phi) (i\partial_\tau \phi + v\partial_x \phi). \quad (3)$$

Here v is the velocity of the edge excitation, which is non-universal, and will depend on the details of the edge confining potential and the Coulomb interaction at the edge. The dimensionless parameter g on the other hand is *universal* and depends only on the quantum Hall state, $|g| = \nu$. As emphasized by Wen the requirement that the Hamiltonian associated with eq.(2) be bounded below requires that v/g be positive. This analysis neglects the effects of a long range Coulomb interaction which are expected to become relevant at very low temperatures (roughly $T < 10mK$ for a Hall bar of length $100\mu\text{m}$ and width $1\mu\text{m}$) [4,15].

It follows from eq.(1) that the one-dimensional electron density along the edge, $\rho(x)$, is given by $\rho = \partial_x \phi / 2\pi$. An expression for the electron creation operator at the edge can be obtained by combining this with the fact that the momentum operator conjugate to ϕ is $\Pi_\phi = \rho/g$. Since adding an extra electron to the edge is equivalent to creating an ‘instanton’ in ϕ , in which ϕ changes by 2π in the region near x , the electron creation operator at the edge is simply

$$\psi(x) \sim \exp \left[2\pi i \int^x \Pi_\phi(x') dx' \right] = e^{i\phi(x)/g}. \quad (4)$$

A ‘vortex’ or Laughlin quasiparticle at the edge is created simply by $e^{i\phi(x)}$, which carries fractional charge ge .

We now suppose that the right-moving and left-moving edges are coupled via a tunneling term, say at $x = 0$. The total action will then have the form, $S_L[\phi_L] + S_R[\phi_R] + V(\phi_L, \phi_R)$. Lacking a specific model we can not say whether quasiparticles or electrons will tunnel more easily. Presumably the fractionally-charged quasiparticles see a lower barrier, but the matrix elements may compensate. Instead, we write down the most general form allowed by symmetry. Taking the weak link region to be at $x = 0$, we have

$$V = \sum_{m=1}^{\infty} v_m \exp [im(\phi_L(x=0) - \phi_R(x=0))] + \text{c.c.}, \quad (5)$$

where the v_m are (complex) tunneling amplitudes. The term $m = 1$ represents the combined amplitudes for a quasi-electron to tunnel from one edge to the other *or* a quasi-hole to tunnel in the opposite direction. These physically distinct processes lead to the same final state and hence add coherently to produce v_1 . The term $m = 1/g$ corresponds to electron tunneling. We have no *a priori* knowledge of the v_m . Fortunately, for $g = 1/3$, all terms except v_1 are irrelevant, having a negative renormalization group eigenvalue, $1 - gm^2$, [7]. Thus at low enough temperatures ($T \ll \Delta$, where $\Delta \sim 1\text{K}$ is the bulk excitation gap) and small enough v_1 , the irrelevant variables $v_m, m > 1$, will flow to zero before $v_{\text{eff}} \sim v_1/T^{1-g}$ has grown large. Thus the RG flow will follow a *universal* trajectory away from the resonance fixed point ($v_1 = 0$) into the insulating fixed point.

At finite temperature the renormalization group flows will be cut off, and the system will end up somewhere along that universal trajectory. From this it follows that in the limit of low temperature, the conductance as a function of the resonance tuning parameter $\delta \sim |v_1|$ and the temperature will obey the scaling form,

$$G(T, \delta) = \tilde{G}_g(c\delta/T^{1-g}), \quad (6)$$

The scaling function $\tilde{G}_g(X)$ is *universal* in the sense that it does not depend on microscopic details, but is a property of the universal trajectory connecting the two fixed points. Since $g = \nu$ in the quantum Hall effect, $\tilde{G}_g$ is completely determined by the theory. The parameter c is a non-universal dimensionful factor which sets the temperature scale. By demanding that the scaling form (6) matches onto the off-resonance conductance, which vanishes as $G(T) \sim T^{2/g-2}$, implies that the tails of the scaling function should decay like $X^{-2/g}$, or X^{-6} for $g = 1/3$.

Though the general properties of the scaling function, such as the temperature dependence of the width and the exponent in the tails are known, the detailed shape of the scaling function has been calculated analytically [7] only for $g = 1/2$. This problem is ideally suited to Monte Carlo simulation, and we have explicitly computed $\tilde{G}(X)$, verified the predicted scaling behavior, and determined the entire scaling function. Following ref. [7] we note that

the action is gaussian in $\phi(x)$ for $x \neq 0$ and so we integrate out all degrees of freedom except $\phi(\tau) \equiv \phi_L(x=0, \tau) - \phi_R(x=0, \tau)$. This gives the action,

$$S = \frac{1}{4\pi|g|} \sum_{i\omega_n} |\omega_n| |\phi(\omega_n)|^2 + v_1 \int_0^\beta d\tau \cos \phi(\tau), \quad (7)$$

where we have retained only the single relevant operator. We have computed the finite frequency ‘two-terminal’ conductance from the Kubo formula,

$$G(\omega_n) = \frac{e^2}{2\pi\hbar} |\omega_n| \langle |\phi(\omega_n)|^2 \rangle. \quad (8)$$

A hard cutoff Λ is introduced by keeping only a finite number of Matsubara frequencies L (typically $L < 100$). We also simulated a dual version of the model in which the tunneling events are represented by a plasma of logarithmically interacting ‘charges’ [7]. Essentially identical results were obtained in the two approaches.

In order to extract information about the temperature dependence of the D.C. conductance, analytic continuation to zero frequency is necessary. Though difficult to do exactly, this may be done with sufficient accuracy by fitting the finite frequency data to a rational function [2/3] Padé form in order to extrapolate to $\omega = 0$.

In order to check the code and to test our analytic continuation procedure, we compare our results to an exact solution which is available for $g = 1/2$ in Fig.(2) The solid line is the exact solution derived in ref. [7], and the data points correspond to Monte Carlo simulations of G as a function of cv_1/T^{1-g} for different system sizes corresponding to $T = \Lambda/(31\pi)$ and $T = \Lambda/(41\pi)$. c is the single non-universal parameter which is adjusted to obtain the fit.

In Fig.(3) we display the results of our Monte Carlo simulation for $g = 1/3$. In this case the tails of the resonance are predicted to decay much faster, like X^{-6} . The data clearly scales. If the coefficient c in eq.(7) is chosen such that the scaling function varies as $\tilde{G}(X) = g(1 - X^2)$ for small X , this simulation allows us to determine that for large X $\tilde{G}(X) = KX^{-6}$ with $K = 2.6 \pm 0.2$.

It should be emphasized that this scaling behavior is to be expected for the fractional quantum Hall effect $\nu = 1/3$, but not for the integer effect $\nu = 1$ or higher-order fractions.

In the integer case the edge state is equivalent to a non-interacting Fermi-liquid, and at low temperatures the resonances should be temperature-independent and Lorentzian. For all higher-order fractions in Laughlin's sequence, $1/\nu$ an odd integer, multi-quasiparticle backscattering processes, v_m in (5) for $m > 1$, are also relevant and grow at low temperatures. Thus in these cases the conductance at the peak of the "resonance" ($v_1 = 0$) will decrease upon cooling, eventually killing completely the resonance in the zero temperature limit. The higher order hierarchical quantum Hall fluids, such as $\nu = 2/3, 2/5, 2/7, \dots$, have more than one branch of edge states [4,16], which complicates the analysis. Resonant tunneling between hierarchical edges will be considered in a subsequent paper.

ACKNOWLEDGMENTS

It is a pleasure to thank V. Chandrasaekar, A. H. MacDonald, F. Milliken, H. Mooij, Mats Wallin and R. Webb for thoughtful discussions. M.P.A.F. is grateful to the Institute for Theoretical Physics in Santa Barbara where part of this work was carried out. The work at Indiana University was supported by NSF DMR-9113911. The work at Pennsylvania was supported by the NSF MRL program, under grant no. DMR91-20668.

REFERENCES

- [1] For a review, see D.V. Averin and K.K. Likharev, in *Mesoscopic Phenomena in Solids*, edited by B. L. Altshuler, P.A. Lee and R.A. Webb (Elsevier, Amsterdam, 1990).
- [2] Y. Meir, N.S. Wingreen and P.A. Lee, *Phys. Rev. Lett.* **66**, 3048 (1991).
- [3] C.L. Kane and M.P.A. Fisher, *Phys. Rev. Lett.* **68**, 1220 (1992).
- [4] X.G. Wen, *Phys. Rev. B* **43**, 11025 (1991); *Phys. Rev. Lett.* **64**, 2206 (1990). X.G. Wen, *Phys. Rev. B* **44** 5708 (1991).
- [5] C. de C. Chamon and X. G. Wen (preprint, 1993).
- [6] A. Furusaki and N. Nagaosa, *Phys. Rev. B* **47**, 3827 (1993).
- [7] C. L. Kane and M. P. A. Fisher, *Phys. Rev. B* **46**, 7268 (1992); *op. cit.*, 15233 (1992).
- [8] These tunneling paths (weak links) may be due to random impurity potentials or an intentionally created quantum dot. If the tunneling occurs through a dot with significant charging energy U , or the tunneling matrix element between the dot from the leads is very small, a more careful analysis is required in order to see how the resonance fixed point is reached. This has been studied in detail in ref. [7].
- [9] Since the Laughlin quasiparticles are vortices, the relative phase for the tunneling on the two weak links is given by the number of enclosed electrons in the area between the two links. This is adjustable by means of the B field and possibly electrostatically.
- [10] Or equivalently the four-terminal conductance G^{13} with the voltage probes on opposite sides of the sample and straddling the weak-link region as shown in Fig.(1).
- [11] J. Frölich and A. Zee, *Nucl. Phys. B* **364**, 517 (1991).
- [12] S. M. Girvin and A. H. MacDonald, *Phys. Rev. Lett.* **58**, 1252 (1987).
- [13] S-C. Zhang, T. H. Hansson and S. Kivelson, *Phys. Rev. Lett.* **62**, 82 (1989).

- [14] D. H. Lee and M. P. A. Fisher, Phys. Rev. Lett. **63**, 903 (1989).
- [15] L.I. Glazman, I.M. Ruzin and B.I. Shklovskii, Phys. Rev. B **45** 8454 (1992).
- [16] A. H. MacDonald, Phys. Rev. Lett. **64**, 222 (1990); M. D. Johnson and A. H. MacDonald, Phys. Rev. Lett. **67**, 2060 (1991).

FIGURES

FIG. 1. Four terminal Hall bar geometry with a narrow constriction between the source (S) and drain (D) formed by a lithographically patterned gate (G). The dotted lines represent two parallel tunneling paths for quasiparticles. Precisely on resonance, there is perfect edge transmission between the source (S) and the drain (D) as indicated by the arrows. Away from resonance the edge channels are reflected at low temperatures.

FIG. 2. Log-log plot comparing resonance scaling function for $g = 1/2$ obtained by Monte Carlo simulation for two different temperatures (squares and hexagons) and the exact solution (solid line). The dashed line is the asymptotic behavior predicted to decay as X^{-4} .

FIG. 3. Log-log plot of resonance scaling function for $\nu = 1/3$ fractional quantum Hall effect obtained by Monte Carlo simulation for two different temperatures. The dashed line is the asymptotic behavior predicted to decay as X^{-6} .